

Addendum to

“Decoupling of pion coupling f_π from quarks at high density in three models, and its possible observational consequences”

Phys. Lett. B 618 (2005) 115

DOI: 10.1016/j.physletb.2005.05.031

Manjari Bagchi, Monika Sinha, Mira Dey, Jishnu Dey

February 2, 2008

In two recent papers the decoupling of the quarks from pions at high density is totally ignored. In the more recent one - Baldo *et al.* [1], the authors **assume** that the f_π remains constant and put the cut off as a variable with increasing density. But from 1985 onward various groups including Dey, Dey and Ghose [2], Dey and Dey [3], Bernard, Meissner and Zahed [4] and Eletsky and Kogan [5] and finally Bagchi *et al.* [7] have shown that in various models including QCD sumrule, the pions decouple from quarks at high density. The values of f_π decreases with increasing density. Baldo *et al.* considers the model of Nambu and Jona-Lasinio which is discussed in the work by Bagchi *et al.* [7].

These papers [2, 3, 4, 5] also point out that the pion coupling to the nucleon $g_{\pi NN}$ survives at high density. This implies that the long range tensor force remains unaltered at high density. Klähn *et al.* [6] have recently pointed out that constraints can be derived for the high-density nuclear equation of state from phenomenology of compact stars and heavy-ion collisions. Their scheme is applied to a set of relativistic EoSs constrained otherwise from nuclear matter saturation properties. The result is that no EoS can satisfy all constraints, but those with density-dependent masses and coupling constants appear most promising. Perhaps a density dependent form for the tensor force will be very useful for deriving a new EoS.

Eletsky and Kogan [5] argued that it is worth asking a question whether other hadronic couplings involving pions share this feature. This may then affect hyperon stars which are neutron stars with presence of strange baryons.

The second work is the more interesting one due to Soni and Bhattacharya [8] where they suggest that the energy of ud matter may be lowered in the presence of pion condensate. They claim this imposes more stringent conditions for the stability of uds matter which has to be more bound than ud matter - while ud matter is less bound than nucleons in Fe^{56} . The conditions should be revised with a density dependent f_π , and the condition will slacken.

References

- [1] M. Baldo, G. F. Burgio, P. Castorina, S. Plumari and D. Zappalà, “Neutron Star stability in the Nambu - Jona-Lasinio model and quark confinement”, hep-ph/0607343.
- [2] J. Dey, M. Dey, P. Ghose, Phys. Lett. B 165 (1985) 74.
- [3] J. Dey and M. Dey, Phys. Lett. B 176 (1986) 469.
- [4] V. Bernard, U-G Meissner, I Zahed, Phys. Rev. D 36 (1987) 819.
- [5] V. L. Eletsky, I. I. Kogan, Phys. Rev. D 49 (1994) 3083.
- [6] T. Klähn, D. Blaschke, S. Typel *et al.* “Constraints on the high-density nuclear equation of state from the phenomenology of compact stars and heavy-ion collisions”, MPG-VT-UR 270/06; nucl-th/0602038.
- [7] M. Bagchi, M. Sinha, M. Dey, J. Dey Phys. Lett. B 618 (2005) 115.
- [8] V. Soni, D. Bhattacharya, Phys. Rev. D 69 (2004) 074001; nucl-th/0602038